INT PUB-06-09
NT@UW-06-11

Dimer scattering in the ε expansion

Gautam Rupak[*]

Institute for Nuclear Theory, University of Washington, Seattle, WA 98195 and

Nuclear Theory Group, University of Washington, Seattle, WA 98195

The atom-dimer and dimer-dimer scattering lengths are analytically calculated in an expansion around four spatial dimensions for fermions with a large 2-body scattering length a. We find the atom-dimer scattering length $a_{ad}/a = \frac{4}{3} - \frac{2}{9}\varepsilon + \mathcal{O}(\varepsilon^2)$ and the dimer-dimer scattering length $a_{dd}/a = \varepsilon - 0.344\varepsilon^2 + \mathcal{O}(\varepsilon^3)$, where $\varepsilon = 4 - d$ and d is the number of spatial dimensions. These ratios $a_{ad}/a = 1.11$ and $a_{dd}/a = 0.656$ at $\varepsilon = 1$ are to be compared with the non-perturbatively calculated numerical results 1.18 and 0.6 respectively. The neutron-deuteron scattering length in the quartet channel $a_{nD} \approx 4.78$ fm is in reasonable agreement with the experimental value $a_{nD}^{\mathrm{exp}} = 6.35 \pm 0.02$ fm, considering 2-body effective range corrections $r_0/a \sim 0.4$ were excluded. The deuteron-deuteron scattering length $a_{DD} \approx 3.15$ fm in the spin-2 channel.

arXiv:nucl-th/0605074v1 29 May 2006

Systems with a large 2-body scattering length a are common in nuclear and atomic physics. Recent experiments near the Feshbach resonance have added new systems to this list [1, 2, 3, 4, 5]. Near the unitarity limit $a \to \pm\infty$, the leading behavior is expected to be universal on dimensional ground. However, theoretical calculations in this interesting regime of large 2-body scattering length are difficult – typically involving non-perturbative resummation of diagrams or non-trivial numerical calculations. In Ref. [6], Nussinov and Nussinov pointed out that in $d = 4$ spatial dimensions one obtains a non-interacting system of dimers. Recently Nishida and Son have developed an $\varepsilon = 4 - d$ expansion around four spatial dimensions [7], allowing them to perturbatively calculate thermodynamic properties for a dilute Fermi gas at the unitarity limit $a \to \pm\infty$.

In this paper, we consider the calculation of the atom-dimer and dimer-dimer scattering amplitude at large but finite 2-body S-wave scattering length a such that the interaction is completely determined by 2-body physics. Besides providing a perturbative method for few-body calculations in the vacuum, this work would allow study of many-body properties at finite a such as the Bose-Einstein condensation (BEC) in atomic systems near the Feshbach resonance. The relevant theory for large scattering length a is described by the following Lagrangian density:

$$\mathcal{L} = \psi^\dagger\left(i\partial_0 + \frac{\nabla^2}{2M}\right)\psi - \frac{c_0}{4}\left(\psi^T\sigma_2\psi\right)^\dagger\left(\psi^T\sigma_2\psi\right)\,, \tag{1}$$

where ψ are spin-$\frac{1}{2}$ fermionic fields and the pauli matrix σ_2 projects the interaction onto the singlet S-wave. An equivalent form for the Lagrangian density after a Hubbard-Stratonovich transformation is

$$\begin{aligned}\mathcal{L} = {}&\psi^\dagger\left(i\partial_0 + \frac{\nabla^2}{2M}\right)\psi + \frac{g}{2}\left[\phi^\dagger\left(\psi^T\sigma_2\psi\right) + h.c.\right] \\ &+ \frac{g^2}{c_0}\phi^\dagger\phi\,.\end{aligned} \tag{2}$$

In $d = 3$ spatial dimensions, it is possible to establish a systematic perturbation in R/a where the range of the interaction $R \ll a$ [8, 9, 10, 11]. This has been quite successful in the 2 and 3-nucleon systems, see for example Refs. [12, 13, 14], where the 2-body scattering length is large. However, it involves resumming certain diagrams to all order in perturbation. In Fig. 1, the atom-atom scattering amplitude is $\mathcal{O}(a)$ and requires resumming an infinite set of diagrams all of which are leading order in the perturbation [8, 9, 10, 11]. The leading order atom-dimer amplitude is again $\mathcal{O}(a)$ and given by an infinite set of diagrams, see Ref. [10]. The dimer-dimer amplitude is also $\mathcal{O}(a)$, and similarly requires resumming the indicated diagrams at leading order [15]. The atom-dimer amplitude for spin-$\frac{1}{2}$ fermions has been non-perturbatively calculated model-independently in effective field theory [10, 12, 16]. For dimer-dimer scattering only a partial resummation of the diagrams has been done in Ref. [17].



FIG. 1: The *leading order* scattering amplitudes for atoms and dimers in $d = 3$ spatial dimensions. Single lines: atoms/fermions, double lines: *fully dressed* dimers/bosons.

Working in $d = 4$ spatial dimensions will avoid the non-perturbative treatment of the dimer mentioned above [6]. We

---

[*] Electronic address: grupak@u.washington.edu

Typeset by REVTEX

will formulate an ε expansion for atom-dimer and dimer-dimer scattering in $d = 4 - \varepsilon$ dimensions [7]. First we define the coupling c_0 in arbitrary d spatial dimensions using the pole in the 2-body scattering amplitude, Fig. 1

$$i\mathcal{T}_{aa} = \frac{-ic_0}{1 + ic_0 L(p_0, \vec{p})} \tag{3}$$

located at the 2-body binding energy $B = 1/(a^2 M)$:

$$\begin{aligned} \frac{1}{c_0} &= -\, iL(p_0, \vec{p})\Big|_{p_0 = -B + p^2/(4M)} \\ &= -\,\frac{M}{(4\pi)^{d/2}}\,(MB)^{\frac{d-2}{2}}\,\Gamma\left(1 - \frac{d}{2}\right), \end{aligned} \tag{4}$$

where

$$\begin{aligned} L(p_0, \vec{p}) &= -\, iM \int \frac{d^d\, q}{(2\pi)^d}\,\frac{1}{q^2 - Mp_0 + \frac{p^2}{4} - i0^+} \\ &= -\, i\frac{M}{(4\pi)^{d/2}}\left(-Mp_0 + \frac{p^2}{4} - i0^+\right)^{\frac{d-2}{2}}\Gamma\left(1 - \frac{d}{2}\right), \end{aligned} \tag{5}$$

and then write the Lagrangian density in terms of the kinetic and interaction piece as

$$\begin{aligned} \mathcal{L} &= \psi^\dagger\left(i\partial_0 + \frac{\nabla^2}{2M}\right)\psi + \phi^\dagger\left(i\partial_0 + \frac{\nabla^2}{4M} + B\right)\phi + \mathcal{L}_I\,, \\ \mathcal{L}_I &= -\,\phi^\dagger\left(i\partial_0 + \frac{\nabla^2}{4M} + B\right)\phi + \frac{g}{2}\left[\phi^\dagger\left(\psi^T\sigma_2\psi\right) + h.c.\right] \\ &\quad + \frac{g^2}{c_0}\phi^\dagger\phi\,. \end{aligned} \tag{6}$$

The original Lagrangian density in Eq. (1) had only one coupling c_0 that was determined by the 2-body scattering length a, the only parameter in the theory. In Eq. (2), after the Hubbard-Stratonovich transformation, we have an additional coupling g. However, its value is arbitrary and only the ratio of the yukawa coupling squared g^2 and the "mass" term g^2/c_0, i.e. c_0, contributes after the dimer wavefunction renormalization. In $d = 3$ dimensions, the wavefunction renormalization for the dimer $Z_d \sim 1/a$ [11]. In $d = 4 - \varepsilon$ dimensions, where we expect a non-interacting theory of dimers [6], the coupling g is chosen such that using the Lagrangian density in Eq. (6) gives $Z_d \sim 1 + \mathcal{O}(\varepsilon)$. This then requires the dimer self-energy $i\Sigma(p_0, \vec{p})$ in Fig. 2 to vanish on-shell and only contribute to Z_d through a derivative term at $\mathcal{O}(\varepsilon)$:

$$\begin{aligned} i\Sigma(p_0, \vec{p}) &\approx \varepsilon\left(p_0 + B - \frac{p^2}{4M}\right) i\frac{\partial\Sigma}{\partial p_0}\Big|_{p_0 = -B + p^2/(4M)}, \\ Z_d &= 1 - \varepsilon\frac{\partial\Sigma}{\partial p_0}\Big|_{p_0 = -B + p^2/(4M)} + \cdots, \end{aligned} \tag{7}$$

where "$\cdots$" indicate terms higher order in the ε expansion.

In $\varepsilon = 4 - d$ dimensions, for Eq. (7) to hold, the coupling g has to be

$$g = \frac{\sqrt{8\pi^2\varepsilon}}{M}\underbrace{M^{\varepsilon/2}}_{\text{arbitrary scale}}, \tag{8}$$



FIG. 2: Dimer wavefunction renormalization. Solid lines: atoms/fermions, dashed lines: dimers/bosons. Unlike in $d = 3$ dimensions, the dimer propagators are not dressed since the dimer-fermion interaction is perturbative around $d = 4$. The "X" represents insertion of ig^2/c_0 and the "square" kinetic energy $-i[p_0 + B - \vec{p}^2/(2M)]$ from $\mathcal{L}_I$ in Eq. (6), on the dimer propagator.

where only the factor $\frac{\sqrt{8\pi^2\varepsilon}}{M}$ is uniquely determined from the dimer wavefunction renormalization. With this choice of the coupling g, it is straightforward to estimate the relative sizes of diagrams. For elastic scattering, diagrams involve even powers of g and we associate a power of ε for every factor of g^2. In effect, the ε expansion is a loop expansion.

For the wavefunction renormalization Z_d, we get:

$$\begin{aligned} i\Sigma(p_0, \vec{p}) &= -\, g^2 L(p_0, \vec{p}) + i\frac{g^2}{c_0} - i\left(p_0 + B - \frac{p^2}{4M}\right), \\ Z_d &\approx 1 + \frac{\varepsilon}{2}\left[\gamma_E - \log(4\pi a^2 M^2)\right], \end{aligned} \tag{9}$$

where $\gamma_E \approx 0.57722$ is the Euler-Mascheroni constant.

The atom-atom scattering amplitude is given by the diagrams in Fig. 3. We find at threshold:

$$\frac{i\mathcal{T}_{aa}}{a^{3-\varepsilon}} = -\, i\frac{8\pi^2}{aM}\varepsilon + i\frac{4\pi^2}{aM}\left[1 - \gamma_E + \log(4\pi)\right]\varepsilon^2 + \cdots, \tag{10}$$

where we divided the amplitude $i\mathcal{T}_{aa}$ by factors $a^{3-\varepsilon}$ to look at dimensionless ratios for convenience. Unlike in $d = 3$ dimensions, the atom-atom scattering amplitude $i\mathcal{T}_{aa}$ scales as $\mathcal{O}(a^2)$ in $d = 4$ dimensions.



FIG. 3: Atom-atom scattering amplitude $\mathcal{T}_{aa}$ in the ε expansion. Diagrams up to the next-to-leading order are shown. Same notation as in Fig. 2.

A non-trivial check of the ε expansion with finite a would be the atom-dimer scattering length. Like the atom-atom scattering amplitude, the atom-dimer amplitude $i\mathcal{T}_{ad}$ also starts at

$\mathcal{O}(\varepsilon)$, see Fig. 4. At $\mathcal{O}(\varepsilon^2)$, there is a contribution from a 1-loop diagram shown in Fig. 4, in addition to the wavefunction renormalization factor Z_d calculated in Eq. (9). The contribution from the tree diagram (including a factor of $a^{\varepsilon-3}$) in Fig. 4 is:

$$-i\frac{g^2a^2MZ_d}{a^{3-\varepsilon}} = -i\frac{8\pi^2}{aM}\varepsilon - i\frac{4\pi^2}{aM}\left[\gamma_E - \log(4\pi)\right]\varepsilon^2 + \cdots . \tag{11}$$

Together with the contribution from the 1-loop diagram, we get at next-to-leading order

$$\frac{i\mathcal{T}_{ad}}{a^{3-\varepsilon}} = -i\frac{8\pi^2}{aM}\varepsilon - i\frac{4\pi^2}{3aM}\left[3\gamma_E - 4 - 3\log(4\pi)\right]\varepsilon^2 . \tag{12}$$

To compare with numerical results calculated in 3 spatial dimensions, we consider the dimensionless ratio

$$\frac{\mathcal{T}_{ad}}{\mathcal{T}_{aa}} = 1 - \frac{\varepsilon}{6} + \mathcal{O}(\varepsilon^2) \approx 0.83\Big|_{\varepsilon=1} . \tag{13}$$

In $d = 3$ spatial dimensions, $\mathcal{T}_{ad}/\mathcal{T}_{aa} = 3a_{ad}/(4a)$. Numerical evaluation by solving the 3-body Faddeev equation [10, 12, 16]

$$\begin{aligned}
\mathcal{T}_{ad} &= \frac{3\pi a}{M} b(3x^2/4-1, x, x)\Big|_{x=0} , \qquad (14)\\
b(\eta, x, z) &= -K(\eta, x, z) - \frac{2}{\pi}\int_0^\infty dy \frac{y^2 b(\eta, x, y)}{y^2 - \frac{4}{3}(\eta+1)} K(\eta, y, z) ,\\
K(\eta, x, z) &= \frac{2}{3}\frac{1+\sqrt{\frac{3}{4}x^2-\eta}}{xz}\log\left[\frac{x^2+z^2+xz-\eta}{x^2+z^2-xz-\eta}\right] ,
\end{aligned}$$

gives $\mathcal{T}_{ad}/\mathcal{T}_{aa} \approx 0.885$ ($a_{ad} \approx 1.18a$). The ε expansion result is perturbatively close to the non-perturbatively calculated numerical value, and it is a tremendous simplification over the non-perturbative calculation in $d = 3$ dimensions.



FIG. 4: Atom-dimer scattering amplitude $\mathcal{T}_{ad}$ in the ε expansion. Diagrams up to the next-to-leading order are shown. Same notation as in Fig. 2.

The discussion can be generalized to certain spin channels in nuclear physics where one has to introduce isospin degrees of freedom to accommodate both the proton and the neutron with similar properties under the strong nuclear interaction. Specifically, the dimer in nuclear physics context is the deuteron, a bound state of a proton and a neutron, in the spin-1 channel. The spin degrees of freedom in the dimer calculation can be thought of as the isospin of nuclear physics, and then the dimer calculation can be applied to deuteron scattering in channels where all the proton and neutron spins are pointing in the same direction. Thus the atom-dimer amplitude is the same as the neutron-deuteron scattering in the S-wave quartet channel with angular momentum $J = 3/2$.

The deuteron binding energy of $B = 2.2246$ MeV corresponds to $a = 1/\sqrt{MB} = 4.32$ fm, ignoring the effective range $r_0 = 1.75$ fm [18]. The standard nuclear physics definition of the scattering length $\hat{a} \approx a + r_0/2 \approx 5.2$ fm. From the atom-dimer calculation, the neutron-deuteron quartet channel scattering length is $a_{nD} \approx 1.11a \approx 4.78$ fm. This is in reasonable agreement with the experimental value $a_{nD}^{\mathrm{exp}} = 6.35 \pm 0.02$ fm [19] considering we neglected $r_0/a \sim 0.4$ effects. For the deuteron, the effective range r_0 corrections are significant even though formally higher order in the perturbation. The so-called Z-parameterization was developed in $d = 3$ dimensions to precisely account for these effects [20].

Finally, the dimer-dimer scattering length is calculated in the ε expansion. The leading contribution to the dimer-dimer scattering is $\mathcal{O}(\varepsilon^2)$ from the 1-loop diagram shown in Fig. 5. We calculate the dimer-dimer scattering amplitude to next-to-leading order where it receives a 2-loop contribution. The 1-loop contribution is:

$$\begin{aligned}
&-i\frac{2M^3Z_d^2g^4}{a^{3-\varepsilon}}\int\frac{d^d\,q}{(2\pi)^d}\frac{1}{[q^2+1/a^2]^3} \qquad (15)\\
&= -i\frac{2M^3Z_d^2g^4}{(4\pi)^{d/2}}\frac{a^{6-d}}{a^{3-\varepsilon}}\frac{\Gamma(3-d/2)}{\Gamma(3)}\\
&= -i\frac{4\pi^2}{aM}\varepsilon^2 - i\frac{2\pi^2}{aM}\left[\gamma_E - \log(4\pi)\right]\varepsilon^3 + \cdots .
\end{aligned}$$

The 2-loop contribution gives

$$\begin{aligned}
&i\frac{2M^5Z_d^2g^6}{a^{3-\varepsilon}}\int\frac{d^d\,q}{(2\pi)^d}\frac{d^d\,l}{(2\pi)^d}\frac{1}{[q^2+1/a^2]^2\,[l^2+1/a^2]^2} \qquad (16)\\
&\qquad\frac{1}{(q+l)^2/4+q^2/2+l^2/2+1/a^2}\\
&= i\frac{16\pi^2}{3aM}\left[1-18\log(3)+14\log(4)\right]\varepsilon^3 + \mathcal{O}(\varepsilon^4) .
\end{aligned}$$

Thus, for the next-to-leading order dimer-dimer amplitude, we get the analytic result:

$$\begin{aligned}
\frac{i\mathcal{T}_{dd}}{a^{3-\varepsilon}} &= -i\frac{4\pi^2}{aM}\varepsilon^2 + i\frac{2\pi^2}{3aM}\left[8 - 3\gamma_E + 224\log(2)\right. \qquad (17)\\
&\qquad \left. -144\log(3) + 3\log(4\pi)\right]\varepsilon^3 ,\\
\frac{\mathcal{T}_{dd}}{\mathcal{T}_{aa}} &= \frac{1}{2}\varepsilon - 0.172\varepsilon^2 + \mathcal{O}(\varepsilon^3) \approx 0.328\Big|_{\varepsilon=1} .
\end{aligned}$$

$\mathcal{T}_{dd}/\mathcal{T}_{aa} = a_{dd}/2a$ in $d = 3$ spatial dimensions and numerical calculation gives $a_{dd} = 0.6a$ [21], a value perturbatively reproduced in the epsilon expansion. From the dimer-dimer calculation, the deuteron-deuteron scattering length in channels with all the proton and neutron spins in the same direction is $a_{DD} \approx 3.15$ fm in the ε expansion, ignoring 2-body effective range r_0 effects.

Systems with unnaturally large 2-body scattering lengths are common in nature and typically necessitate non-perturbative calculations that are computationally difficult and expensive. Study and development of perturbative techniques



FIG. 5: Dimer-dimer scattering amplitude $\mathcal{T}_{dd}$ in the ε expansion. Diagrams up to the next-to-leading order are shown. Same notation as in Fig. 2.

are always useful. Many times perturbative calculations can be done analytically that allow insights otherwise not possible in a numerical calculation. In this paper, we considered a fermi system with a large but finite 2-body scattering length a in an expansion around $d = 4$ spatial dimensions. The effect of the 2-body scattering length was systematically incorporated and the atom-dimer and dimer-dimer scattering length was calculated to next-to-leading order in perturbation. The results seem to converge to the non-perturbatively calculated numerical solutions at this order of the calculation, and the ε expansion should be an useful tool for studying strongly interacting systems with large 2-body scattering length a. For example, the current work would be important for research in the BEC-BCS crossover region in atomic systems as one varies the scattering length a across the Feshbach resonance. As shown, the dimer calculation can be generalized to deuterons in nuclear physics in certain spin channels. However, it is not clear from this calculation how the Efimov effect [22] in bosons and fermions with isospin (more than one fermion species as in the proton-neutron system) associated with non-perturbative renormalization [23] emerges in the ε expansion. It could be that the Efimov effect cannot be described by an analytic continuation from $d = 4$ spatial dimensions. It is known that the Efimov effect occur only in spatial dimensions $2.3 < d < 3.8$, with $d = 3$ being the only integer dimension [24].

Acknowledgments

The author thanks D. T. Son for helpful discussions, and P. F. Bedaque and D. B. Kaplan for comments on the manuscript. This work was supported in part by DOE grants DE-FG02-00ER41132, DE-FC02-01ER41187 and DE-FG02-97ER41014.

---

[1] K. M. O'Hara, S. L. Hemmer, M. E. Gehm, S. R. Granade and J. E. Thomas, *Science* **298**, 2179 (2002); M. E. Gehm, S. L. Hemmer, S. R. Granade, K. M. O'Hara and J. E. Thomas, *Phys. Rev.* A **68**, 011401(R) (2003).

[2] T. Bourdel, J. Cubizolles, L. Khaykovich, K. M. F. Magalhães, S. J. J. M. F. Kokkelmans, G. V. Shlyapnikov and C. Salomon, *Phys. Rev. Lett.* **91**, 020402 (2003).

[3] K. Dieckmann, C. A. Stan, S. Gupta, Z. Hadzibabic, C. H. Schunck and W. Ketterle, *Phys. Rev. Lett.* **89**, 203201 (2002).

[4] C. A. Regal, C. Ticknor, J. L. Bohn and D. S. Jin, *Nature* **424**, 47 (2003); M. Greiner, C. A. Regal and D. S. Jin, *Nature* **426**, 537 (2003).

[5] S. Jochim, M. Bartenstein, A. Altmeyer, G. Hendl, S. Riedl, C. Chin, J. Hecker Denschlag, and R. Grimm, *Science* **302**, 2101 (2003).

[6] Z. Nussinov and S. Nussinov, cond-mat/0410597.

[7] Y. Nishida and D. T. Son, cond-mat/0604500.

[8] D. B. Kaplan, M. J. Savage and M. B. Wise, *Phys. Lett.* **B424**, 390 (1998).

[9] U. van Kolck, Nucl. Phys. **A645**, 273 (1999).

[10] P. F. Bedaque and U. van Kolck, *Phys. Lett.* **B428**, 221 (1998).

[11] J.-W. Chen, G. Rupak and M. J. Savage, *Nucl. Phys.* **A653**, 386 (1999).

[12] P. F. Bedaque, H. W. Hammer and U. van Kolck, *Phys. Rev.* **C58**, 641 (1998).

[13] M. Butler, J.-W. Chen and X. Kong, *Phys. Rev.* **C63**, 035501 (2001).

[14] G. Rupak, *Nucl. Phys.* **A678**, 405 (2000).

[15] T. Luu and G. Rupak, unpublished.

[16] G. Rupak, nucl-th/0604053.

[17] P. Pieri and G. C. Strinati, *Phys. Rev.* B **61**, 15370 (2000).

[18] J. J. de Swart, C. P. F. Terheggen and V. G. J. Stoks, nucl-th/9509032.

[19] W. Dilg, L. Koester and W. Nistler, *Phys. Lett.* **B36**, 208 (1971).

[20] D. R. Phillips, G. Rupak and M. J. Savage, Phys. Lett. **B473**, 209 (2000).

[21] D. S. Petrov, C. Salomon and G. V. Shlyapnikov, *Phys. Rev.* A **71**, 012708 (2005).

[22] V. N. Efimov, *Sov. J. Nucl. Phys.* **12**, 589 (1971); *Phys. Rev.* C **47**, 1876 (1993).

[23] P. F. Bedaque, H. W. Hammer and U. van Kolck, *Phys. Rev. Lett.* **82**, 463 (1999).

[24] E. Nielsen, D. V. Fedorov, A. S. Jenson and E. Garrido, *Phys. Rev.* **347**, 373 (2001).